arXiv:1106.2348v2 [math.AC] 15 Jun 2011

LINEAR QUOTIENTS OF SQUARE OF THE EDGE IDEAL OF THE ANTICYCLE

A. H. HOEFEL AND G. WHIELDON

Abstract. Let G be a graph with chordal complement and $I(G)$ its edge ideal. From work of Herzog, Hibi, and Zheng, it is known that $I(G)$ has linear quotients and all of its power have linear resolutions. For edge ideals $I(G)$ arising from graphs which do not have chordal complements, exact conditions on their powers possessing linear resolutions or linear quotients are harder to find. We provide here an explicit linear quotients ordering for all powers of the edge ideal of the antipath and a linear quotients ordering on the second power $I(A_n)^2$ of the edge ideal of the anticycle A_n. This linear quotients ordering on $I(A_n)^2$ recovers a prior result of Nevo that $I(A_n)^2$ has a linear resolution.

1. Introduction and Background

Let G be a simple graph on n vertices, and $I(G)$ its edge ideal, i.e., a squarefree monomial ideal in $R = \Bbbk[x_1, \ldots, x_n]$ with monomial generators x_ix_j corresponding to each edge $\{i,j\} \in G$. Such ideals have been extensively studied in such papers as [HVT07], [HVT08], [Nev11], [Vil90], and more recently, [MV10]. A goal of much recent research has been to classify behavior of the resolutions of such ideals $I(G)$ and that of their powers in terms of combinatorial data of G. We provide here an explicit proof that the second power of the edge ideal of the anticycle has not just a linear resolution, but also linear quotients.

In the course the proof, we additionally demonstrate that all powers $I(P_n^c)^k$ of the edge ideal of the antipath have linear quotients.

Definition 1.1. Let G be a simple graph on n vertices. Then the *edge ideal of G* is the squarefree monomial ideal $I(G)$ given by

$$I(G) = (x_ix_j : \{i,j\} \in G).$$

We say that a graph G has property P if its edge ideal $I(G)$ has such a property; e.g., G is Gorenstein if $I(G)$ is Gorenstein, G is linear if $I(G)$ has a linear resolution, etc. In particular, we will say a graph G has linear quotients if its edge ideal $I(G)$ has linear quotients:

Definition 1.2. Let I be a homogeneous ideal. We say that I *has linear quotients* if there exists some ordering of the generators of $I = (m_1, m_2, \ldots, m_r)$ such that for all $i > 1$,

$$((m_1, \ldots, m_{i-1}) : (m_i)) = (x_{k_1}, \ldots, x_{k_s})$$

for some variables $x_{k_1}, \ldots, x_{k_s}$. We say that such an ordering $(m_1, m_2, \ldots, m_r)$ is a *linear quotients ordering of I*.

The first author's research is supported by NSERC and Killam scholarships.

2000 *Mathematics Subject Classification.* 13D02.

Key words and phrases. Edge ideal, anticycle, linear quotients, powers of ideals.

For two monomials m and m' we define $m' : m$ to be the monomial $\frac{m'}{\gcd(m,m')}$. Given monomials $m_1, \ldots, m_i$, the colon ideal $(m_1, \ldots, m_{i-1}) : (m_i)$ can be computed as

$$(m_1, \ldots, m_{i-1}) : (m_i) = (m_1 : m_i, \ldots, m_{i-1} : m_i).$$

Thus, in order to show that a monomial ideal $I = (m_1, \ldots, m_r)$ has linear quotients, it suffices to show that for each pair of monomials m_i and m_j with $j < i$ that there exists another monomial m_k with $k < i$ with

$$m_k : m_i = x_l \text{ for some } l \qquad \text{and} \qquad x_l \text{ divides } m_j : m_i.$$

The *graded Betti numbers* of a homogeneous ideal I are given by $\beta_{i,j}(I) = \dim_{\Bbbk} \operatorname{Tor}_i(I, \Bbbk)_j$. The graded Betti numbers also correspond to the ranks of the free modules in a minimal free resolution of I. We say an ideal I which is generated in degree d has a *linear resolution* if $\beta_{i,j}(I) = 0$ for $j \neq i + d$. Ideals with linear quotients also have linear resolutions.

Providing a linear quotients ordering is one technique for proving that an ideal has a linear resolution, often with combinatorial significance in the case of monomial ideals. In the case of squarefree monomial ideal, an ideal I having linear quotients is equivalent to its Alexander dual $I^\vee$ having a shelling order on its facets. For non-squarefree monomial ideals, a linear quotient orderingscan be viewed as giving a shelling order on the Alexander dual of its polarization.

Interest in powers of the anticycle partially draws from a result of Herzog, Hibi and Zheng [HHZ04] which states the following:

Theorem 1.3 (Herzog, Hibi, Zheng)**.** *Let I be a quadratic monomial ideal of the polynomial ring. The following are equivalent:*

1. *I has a linear resolution,*
2. *I has linear quotients,*
3. *I^k has a linear resolution for all $k \geq 1$.*

For edge ideals, Fröberg showed that $I(G)$ has a linear resolution if and only if the complement of G is chordal [Frö90].

Conspicuously missing from the above theorem is the statement that all powers of a quadratic monomial ideal I with linear resolution must have linear quotients. In fact, this is not known. There are numerous examples of non-quadratic monomial ideals possessing a linear resolution, or even linear quotients, whose powers do not. In [Con06], Conca provides a example generated in degree 3 which is not dependent on the characteristic of the field $\Bbbk$.

It would be of interest to construct linear quotients of powers of quadratic monomial ideals with the aim of extending Herzog, Hibi and Zheng's theorem. Alternately, as no counterexamples are known, the construction of a quadratic monomial ideal I with a linear resolution but some power k with no linear quotients ordering on the generators of I^k would be of combinatorial interest.

Our work on the second power of the anticycle was also inspired by a second thread of research. Francisco, Hà and Van Tuyl first investigated graphs G where $I(G)^k$ has a linear resolution for each $k \geq 2$.

From Fröberg and Herzog, Hibi and Zheng's results, we see that chordal graphs have this property. More generally, it has been shown by Francisco, Hà and Van Tuyl that if some power of $I(G)$ has a linear resolution, then the complement of G cannot contain any induced four cycles. Their proof was recorded in [NP09].

Inspired by these results, Peeva and Nevo constructed an example of a graph G with no four cycle in its complement and where $I(G)^2$ does not have a linear resolution. Peeva and Nevo have conjectured that their example works only because $I(G)$ has Castelnuovo-Mumford regularity four and that every successive power of an edge ideal should get strictly closer to a linear resolution. See [NP09] for a more precise statement.

Nevo has also shown that claw-free graphs with no four cycles in their complements have regularity at most three and their second powers have linear resolutions [Nev11]. Anticycles on more than four vertices meet these criteria and so, it follows that their second powers have linear resolutions. Here we demonstrate that the square of the edge ideal of the anticycle has linear quotients, recovering this result.

2. Cycles, Anticycles, and Antipaths

We first describe the edge ideal of the anticycle and partition pairs of its edges into several natural classes. Next, we provide a linear quotients ordering on these classes relative to the previous generators.

The *complement* of a graph G is the graph on the vertices of G containing all edges that are not in G. We use G^c to denote the complement graph.

Definition 2.1. Let C_n be the cycle graph on n vertices, i.e. the graph consisting of one cycle of length n on these vertices with no chords. The *anticycle graph* A_n is the complement graph of C_n, i.e., $A_n = C_n^c$.

Definition 2.2. The *antipath* P_n^c is the graph on n vertices containing of all edges in the complement of a path P_n of length $n-1$. We depict the antipath in the figure below.





Producing a linear quotients ordering for graphs with chordal complements is always possible and all of their powers have linear resolutions, as given in Theorem 3.2 in [HHZ04]. However, most naive orderings on the generators of higher powers of $I(G)$ fail to produce linear quotients for G with chordal complements.

Example 2.3. Let $R = \mathbb{k}[x_1, \dots, x_6]$ and let $I = I(A_n)^2$ be the square of the edge ideal of the anticycle on 6 vertices in R. Its generators, written in lex order, are given by:

$$\begin{aligned}
&x_1^2x_3^2, x_1^2x_3x_4, x_1^2x_3x_5, x_1^2x_4^2, x_1^2x_4x_5, x_1^2x_5^2, x_1x_2x_3x_4, x_1x_2x_3x_5, x_1x_2x_3x_6,\\
&x_1x_2x_4^2, x_1x_2x_4x_5, x_1x_2x_4x_6, x_1x_2x_5^2, x_1x_2x_5x_6, x_1x_3^2x_5, x_1x_3^2x_6, x_1x_3x_4x_5,\\
&x_1x_3x_4x_6, x_1x_3x_5^2, x_1x_3x_5x_6, x_1x_4^2x_6, x_1x_4x_5x_6, x_2^2x_4^2, x_2^2x_4x_5, x_2^2x_4x_6, x_2^2x_5^2,\\
&x_2^2x_5x_6, x_2^2x_6^2, x_2x_3x_4x_5, x_2x_3x_4x_6, x_2x_3x_5^2, x_2x_3x_5x_6, x_2x_3x_6^2, x_2x_4^2x_6,\\
&x_2x_4x_5x_6, x_2x_4x_6^2, x_3^2x_5^2, x_3^2x_5x_6, x_3^2x_6^2, x_3x_4x_5x_6, x_3x_4x_6^2, x_4^2x_6^2.
\end{aligned}$$

This ordering *fails* to be a linear quotients ordering. Let m_i be the i^{th} monomial in the ordering above, and let I_i denote the ideal generated by the first $i-1$ monomials in the ordering. Setting $Q_i = I_i : (m_i)$, we see that

$$\begin{aligned}
Q_9 &= (x_1^2x_3^2, x_1^2x_3x_4, x_1x_2x_3x_4, x_1^2x_4^2, x_1x_2x_4^2, x_2^2x_4^2, x_1^2x_3x_5, x_1x_2x_3x_5) : (x_1x_2x_3x_6)\\
&= (x_4, x_5, x_1x_3)
\end{aligned}$$

is not generated by variables, hence the lex ordering fails to give us linear quotients. Similarly, with reverse lex, we have the following ordered generating set:

$$\begin{aligned}
&x_1^2x_3^2, x_1^2x_3x_4, x_1x_2x_3x_4, x_1^2x_4^2, x_1x_2x_4^2, x_2^2x_4^2, x_1^2x_3x_5, x_1x_2x_3x_5, x_1x_3^2x_5,\\
&x_1^2x_4x_5, x_1x_2x_4x_5, x_2^2x_4x_5, x_1x_3x_4x_5, x_2x_3x_4x_5, x_1^2x_5^2, x_1x_2x_5^2, x_2^2x_5^2, x_1x_3x_5^2,\\
&x_2x_3x_5^2, x_3^2x_5^2, x_1x_2x_3x_6, x_1x_3^2x_6, x_1x_2x_4x_6, x_2^2x_4x_6, x_1x_3x_4x_6, x_2x_3x_4x_6,\\
&x_1x_4^2x_6, x_2x_4^2x_6, x_1x_2x_5x_6, x_2^2x_5x_6, x_1x_3x_5x_6, x_2x_3x_5x_6, x_3^2x_5x_6, x_1x_4x_5x_6,\\
&x_2x_4x_5x_6, x_3x_4x_5x_6, x_2^2x_6^2, x_2x_3x_6^2, x_3^2x_6^2, x_2x_4x_6^2, x_3x_4x_6^2, x_4^2x_6^2.
\end{aligned}$$

This fails to have linear quotients at $Q_{21} = I_{21} : (x_1x_2x_3x_6) = (x_4, x_5, x_1x_3)$. Using a monomial ordering on the generators of I does not appear to ever produce a linear quotients ordering on the generators of $I(A_n)^2$.

This appears to be true more generally – while all higher powers of edge ideals with linear quotients appear to have linear quotients as well, these linear quotients orders almost never arise from a monomial term ordering.

3. ANTIPATH LINEAR QUOTIENTS

Throughout this section we will use $H = P_n^c$ to denote the antipath on n vertices. The first stage in our linear quotients ordering is to show that the square of the antipath has linear quotients with respect to the lex order. As the complement of the antipath is a chordal graph, it is known that $I(H)$ has a linear resolution via Fröberg's Theorem [Frö90]. Furthermore, as $I(H)$ has a linear resolution and is generated in degree 2, it is known to have a linear quotient ordering and linear resolutions of all of its powers [HHZ04]. However, a linear resolution of its second power does not guarantee a linear quotients ordering of $I(H)^k$, which we provide explicitly here.

Proposition 3.1. The k^{th} power $I(H)^k$ of the edge ideal of the antipath H has linear quotients, under the lex ordering of the generators.

We begin with some notation and a lemma.

Given any k edges $e_1, \ldots, e_k$ in a graph G, we will often abuse notation and write $m = e_1 e_2 \cdots e_k$ for the monomial

$$m = \prod_{r=1}^{k} x_{i_r} x_{j_r}$$

where $e_r = \{x_{i_r}, x_{j_r}\}$. When a monomial m is of this form, we say m is the *product of k edges* of G.

Example 3.2. Let G be the complete graph on six vertices $\{x, y, z, w, s, t\}$ seen below.



Then the monomial $m = xyzwst \in I(G)^3$ comes from any three edges with each vertex appearing in a unique edge exactly once.







So $m = e_1 e_2 e_3$ for the labeled edge sets in any of the diagrams above.

Lemma 3.3. The ideal $I(H)^k$ is given by all monomials of degree $2k$ of the form

$$\begin{aligned} I(H)^k = (&x_{i_1} x_{i_2} \cdots x_{i_k} x_{j_1} x_{j_2} \cdots x_{j_k} : \\ &i_1 \leq i_2 \leq \cdots \leq i_k \leq j_1 \leq j_2 \leq \cdots j_k \text{ and } i_r + 2 \leq j_r \text{ for all } r). \end{aligned}$$

Equivalently, every minimal monomial generator $m \in I(H)^k$ can be written as a product of k edges $m = e_1 \cdots e_k$ where $e_r = \{x_{i_r}, x_{j_r}\}$ and

$$i_1 \leq i_2 \leq \cdots \leq i_k \leq j_1 \leq j_2 \leq \cdots \leq j_k.$$

Proof. Any monomial m of degree $2k$ can be written as

$$m = x_{i_1} \cdots x_{i_k} x_{j_1} \cdots x_{j_k}$$

with $i_1 \leq \cdots \leq i_k \leq j_1 \leq \cdots \leq j_k$. Let m be a minimal generator of $I(H)^k$ and write m as above. Assume for a contradiction that there is an index r with $i_r + 2 > j_r$. Since the indices of m have been written in ascending order, we know that

$$\{i_r, i_{r+1}, \ldots, i_k, j_1, \ldots, j_r\} \subseteq \{i_r, i_r + 1\}.$$

Let m' be the degree $k+1$ monomial $m' = x_{i_r} \cdots x_{i_k} x_{j_1} \cdots x_{j_r}$ which divides m. The support of m' is contained in $\{x_{i_r}, x_{i_r+1}\}$ but there are are no edges in the antipath between x_{i_r} and x_{i_r+1}. Thus, m' contains no edge as a factor. However, as m is a product of k edges, every degree $k+1$ factor of m must contain at least one edge. This is contradicted by our construction of m', and so we must have $i_r + 2 \leq j_r$ for each r. □

We now return to the proof of Proposition 3.1.

Proof of Proposition 3.1. From Lemma 3.3, we have that

$$\begin{aligned} I(H)^k = (&x_{i_1}x_{i_2}\cdots x_{i_k}x_{j_1}x_{j_2}\cdots x_{j_k} : \\ &\qquad i_1 \leq i_2 \leq \cdots \leq i_k \leq j_1 \leq j_2 \leq \cdots j_k \text{ and } i_r + 2 \leq j_r \text{ for all } r). \end{aligned}$$

Any pair of monomial generators m and m' of $I(H)^k$ will be of the forms:

$$\begin{aligned} m &= x_{i_1}x_{i_2}\cdots x_{i_k}x_{j_1}x_{j_2}\cdots x_{j_k} = e_1e_2\cdots e_k \\ m' &= x_{i'_1}x_{i'_2}\cdots x_{i'_k}x_{j'_1}x_{j'_2}\cdots x_{j'_k} = e'_1e'_2\cdots e'_k \end{aligned}$$

with indices i_r, i'_r, j_r, j'_r all satisfying the inequalities above and for edges $e_r = \{x_{i_r}, x_{j_r}\}$ and $e'_r = \{x_{i'_r}, x_{j'_r}\}$ of H. We show for every such pair of monomials with $m' >_{\text{lex}} m$ that $m' : m$ will be divisible by some $x_i = m'' : m$ for some $m'' >_{\text{lex}} m$.

Case 1: Monomials m and m' differ first at some x_{i_r}. Assume i_r is the first index at which m and m' differ; i.e., $i_s = i'_s$ for all $s < r$ and $i'_r < i_r$.

Let $m'' = \dfrac{x_{i'_r}}{x_{i_r}}m$. This is certainly a monomial of the appropriate degree which is lex earlier than m. To show that $m'' \in I(H)^k$, we note that as $i'_r < i_r < j_r - 2$, we have an edge $\varepsilon_r = \{x_{i'_r}, x_{j_r}\} \in H$. Thus

$$m'' = e_1\cdots e_{r-1}\varepsilon_r e_{r+1}\cdots e_k \in I(H)^k.$$

As $m'' : m = x_{i'_r}$ and $x_{i'_r}$ divides $m' : m$, we either had $m'' = m'$ (in which case we satisfy the first condition above) or $m'' \neq m'$ and this colon satisfies the second condition above.

Case 2: Monomials m and m' differ first at some x_{j_r}. Assume that m and m' do not differ in the x_{i_s}; i.e., $i_s = i'_s$ for all $s = 1, \ldots, k$. Let j_r be the first index where m and m' differ. That is, $j_s = j'_s$ for all $s < r$ and $j'_r < j_r$. So

$$\begin{aligned} m &= x_{i_1}\cdots x_{i_k}x_{j_1}\cdots x_{j_{r-1}}x_{j_r}x_{j_{r+1}}\cdots x_{j_k} = e_1e_2\cdots e_{r-1}e_re_{r+1}\cdots e_k \\ m' &= x_{i_1}\cdots x_{i_k}x_{j_1}\cdots x_{j_{r-1}}x_{j'_r}x_{j'_{r+1}}\cdots x_{j'_k} = e_1e_2\cdots e_{r-1}e'_re'_{r+1}\cdots e'_k. \end{aligned}$$

Choosing

$$\begin{aligned} m'' &= x_{i_1}\cdots x_{i_k}x_{j_1}\cdots x_{j_{r-1}}x_{j'_r}x_{j_{r+1}}\cdots x_{j_k} \\ &= e_1e_2\cdots e_{r-1}e'_re_{r+1}\cdots e_r, \end{aligned}$$

we note that as $e'_r = \{x_{i_r}, x_{j'_r}\} \in H$, we have $m'' \in I(H)^k$. This is a lex earlier monomial in $I(H)^k$. So $m'' : m = x_{j'_r}$ which divides $m' : m$. □

4. Linear Quotient Ordering of Anticycle

The proof that the square of the edge ideal of the antipath has linear quotients is the first step in constructing a linear quotients ordering of the generators of the anticycle. With this in hand, we now show that the following ordering on the generators of the square of the edge ideal of the anticycle gives us linear quotients. For the remainder of this note, we let G be the anticycle graph and let H be the antipath obtained by deleting some vertex of G.

Remark 4.1. We will label the vertices in G as follows. Let x be the vertex we delete to obtain H, and let z_1 and z_2 the two non-adjacent vertices in G (so the two neighbors of x in the cycle itself). Finally, let $y_1, \dots, y_n$ be all the remainging vertices in order, so that y_1 is not adjacent to z_1 and y_n is not adjacent to z_n. Note that each y_i is adjacent to x. Thus, for this section, we assume that G has $n + 3$ vertices. See the figure below.



Theorem 4.2. Let G be the $(n + 3)$-anticycle graph, labeled as in the picture above, with $n \geq 2$. Let $H = G \setminus \{x\}$ be the induced graph away from x. Let $J = I(H)$ be the edge ideal of H and let $K = I(G \setminus H) = (xy_i : i = 1, \dots, n)$ be the edge ideal on the edges not in H.

Then the edge ideal $I(G)$ has a linear quotients given by the following ordering of its monoimal generators (monomials occurring earlier in this list appear earlier in the order):

(1) $m \in J^2$ ordered via the lex ordering with $z_1 < y_1 < y_2 < \cdots < y_n < z_2$
(2) $m \in J \cdot K$
  (a) $m = xy_iz_1z_2$, $i = 1, \dots, n$,
  (b) $m = xy_iy_jz_2$, $i \leq j$, ordered via lex with $y_1 > y_2 > \cdots > y_n$, excluding nongenerator $xy_n^2z_2$,
  (c) $m = xy_iy_jz_1$, $i \leq j$, ordered via lex with $y_1 < y_2 < \cdots < y_n$, excluding nongenerator $xy_1^2z_1$, and
  (d) $m = xy_iy_jy_k$, $i \leq j \leq k$, ordered via lex with $y_1 > y_2 > \cdots > y_n$.
(3) $m \in K^2$.
  (a) $m = x^2y_iy_j$ ordered via lex excluding $x^2y_1^2$ with $y_1 < y_2 < \cdots < y_n$
  (b) $m = x^2y_1^2$.





Before giving the proof, we provide a specific example of the ordering of $I(G)^2$ for the antipath G on 6 vertices.

Example 4.3. Let $n = 3$ so we have the anticycle graph G on vertices $\{x, z_1, y_1, y_2, y_3, z_2\}$.



Our two subgraphs H and $G \setminus H$ will be as below.





The linear quotients ordering from Theorem 4.2 on the generators of $I(G)^2$ is given here by

$$\begin{aligned} I(G)^2 = \quad & (z_1^2y_2^2, z_1^2y_2y_3, z_1^2y_2z_2, z_1^2y_3^2, z_1^2y_3z_2, z_1^2z_2^2, z_1y_1y_2y_3, z_1y_1y_2z_2, \\ & z_1y_1y_3^2, z_1y_1y_3z_2, z_1y_1z_2^2, z_1y_2^2z_2, z_1y_2y_3z_2, z_1y_2z_2^2, y_1^2y_3^2, \\ & y_1^2y_3z_2, y_1^2z_2^2, y_1y_2y_3z_2, y_1y_2z_2^2, y_2^2z_2^2)^{(1)} \\ & + (xz_1y_1z_2, xz_1y_2z_2, xz_1y_3z_2)^{(2a)} \\ & + (xy_1^2z_2, xy_1y_2z_2, xy_1y_3z_2, xy_2^2z_2, xy_2y_3z_2)^{(2b)} \\ & + (xz_1y_3^2, xz_1y_2y_3, xz_1y_1y_3, xz_1y_2^2, xz_1y_1y_2)^{(2c)} \\ & + (xy_1^2y_3, xy_1y_2y_3, xy_1y_3^2)^{(2d)} \\ & + (x^2y_1y_2, x^2y_1y_3, x^2y_2^2, x^2y_2y_3, x^2y_3^2)^{(\mathrm{3a})} \\ & + (x^2y_1^2)^{(\mathrm{3b})}. \end{aligned}$$

4.1. **Proof of Theorem 4.2.**

Proof of Theorem 4.2. The generators of $I(G)^2$ fall into three main cases, with the second case split up into four subcases and the third case placing the first lex ordered generator at the very end. We will address each case separately.

Note 4.4. Let $I_M = \left(I(G)^2\right)_M$ denote the ideal generated by all monomials in the linear quotients ordering before adding M, a minimal generator of $I(G)^2$. In general, we will use Q_M to denote the colon ideal

$$Q_M = I_M : (M),$$

though we will often omit the subscript if the stage in the ordering is clear. We show here for all monomial generators M in the quotients ordering that

$$Q_M = (x_{i_1}, x_{i_2}, \dots, x_{i_k})$$

for some variables $x_{i_1}, x_{i_2}, \dots, x_{i_k} \in \{x, z_1, z_2, y_1, y_2, \dots, y_n\} = V$.

Let V_M denote the variables generating Q_M, or as above, $V_M = \{x_{i_1}, x_{i_2}, \dots, x_{i_k}\}$ and let $W_M = V \setminus V_M$.

The general technique used begins with generating $x_i \in V_M$ explicitly via exhibition of a monomial generator $m' \in I_M$ such that

$$m' : M = x_i.$$

After finding our expected V_M, we note that any remaining minimal monomial generators m of Q_M which are not variables, i.e. not in a linear generator of the ideal (V_M), must have their support, $\operatorname{supp}(m) \in W_M$.

We then show that any generators $m' \in I(G)^2$ which would give us

$$m' : M = m \in (W_M)$$

must either have $m \in (V_M)$ (and hence a contradiction, as such a generator cannot be minimal in Q_M) or could only come from a monomial m' occurring after M in the linear quotients ordering (and hence another contradiction, as $m \notin Q_M$.) For consistency, we will always use M, m and m' in the same roles throughout the proof.

4.1.1. *Stage (1):* Note that $I(H)$ is the antipath graph of the path $\{z_1 \sim y_1 \sim y_2 \sim \cdots \sim y_n \sim z_2\}$, so the ordering of J^2 given in (1) is a linear quotients ordering by Proposition 3.1.

4.1.2. *Stage (2a):* We now move on to generators in $(2a)$ and show that after adding through the $(i-1)^{\text{st}}$ term in $(2a)$, we have linear quotients when we colon this ideal against our i^{th} term, $M = z_1 z_2 x y_i$. Let Q be this colon ideal,

$$\begin{aligned} Q &= I_{z_1 z_2 x y_i} : (z_1 z_2 x y_i) \\ &= (J^2 + (z_1 z_2 x y_j \mid 1 \le j \le i-1)) : (z_1 z_2 x y_i). \end{aligned}$$

Note that the following inclusions hold, via the elements noted on the right.

- $Q \supseteq (y_j \mid j \neq i)$ as $y_j = z_1 z_2 y_j y_i : z_1 z_2 x y_i$.
- $Q \supseteq (z_1)$ when $i \neq 1$ as $z_1 = z_1 z_2 z_1 y_i : z_1 z_2 x y_i$.
- $Q \supseteq (z_2)$ when $i \neq n$ as $z_2 = z_1 z_2 z_2 y_i : z_1 z_2 x y_i$.
- $Q \supseteq (y_i)$ when $i \notin \{1, n\}$ as $y_i = y_i^2 z_1 z_2 : z_1 z_2 x y_i$.

Assume $m \in Q$ is a minimal monomial generator of Q that is not linear, i.e. $m = m' : z_1 z_2 x y_i$ for some m' appearing in the ordering earlier than $z_1 z_2 x y_i$. As m is minimal, its support cannot contain any of the variables in Q and therefore

$$\operatorname{supp}(m) \subseteq \begin{cases} \{x\} & i = 2, \dots, n-1, \\ \{x, z_1, y_1\} & i = 1, \\ \{x, z_2, y_n\} & i = n. \end{cases}$$

In the first of these cases, we note that if $x | m$ then $x^2 | m'$. As this does not happen for any m' before $z_1 z_2 x y_i$, the only cases we need to consider are $i = 1$ and $i = n$. In both of these cases we can assume that x does not divide m.

Case $(i = 1)$: In this case, we are adding the generator $z_1z_2xy_1$ to J^2, our edge ideal of the antipath, i.e. $Q = J^2 : z_1z_2xy_1$. Note that $Q \supseteq (y_2, \dots, y_n, z_2)$. Hence, if we have a minimal monomial generator $m \in Q$ which is not linear, its support must be contained in $\{z_1, y_1\}$.

If $z_1|m$ then $z_1^2|m'$ so m' must be of the form $z_1^2y_jy_k$ with $j, k > 1$. However, we then have $m' : z_1z_2xy_1 = z_1y_jy_k$ which cannot be a minimal generator of Q, as both $y_j, y_k \in Q$.

If $y_1|m$ then $y_1^2|m'$ so m' must be of the form $y_1^2y_jz_2$ (for $j > 2$) or $y_1^2y_jy_k$ (for $j, k > 2$) or $y_1^2z_2^2$. In these three cases the m' are y_1y_j, $y_1y_jy_k$, and y_1z_2 respectively. However each of these are not minimal, from $y_j, z_2 \in Q$ for $j > 2$.

Case $(i = n)$: Now we are adding the final generator $z_1z_2xy_n$ to the ideal

$$I_{z_1z_2xy_n} = J^2 + (z_1z_2xy_i : 1 \le i \le n - 1).$$

For this, we have $Q = (J^2 + (z_1z_2xy_j \mid 1 \le j \le n - 1)) : (z_1z_2xy_n)$ which satisfies $Q \supseteq (y_1, \dots, y_{n-1}, z_1)$. In this case, if we have a minimal monomial generator $m \in Q$ which is not linear, its support must be contained in $\{z_2, y_n\}$.

If $z_2|m$ then $z_2^2|m'$. The only such $m' \in I_{z_1z_2zy_n}$ must be of the form $z_2^2y_jy_k$ with $j, k < n$. However, we then have $m' : z_1z_2xy_1 = z_2y_jy_k$ which is not a minimal generator as $y_j, y_k \in Q$.

Similarly, if $y_n|m$ then $y_n^2|m'$. All such $m' \in I_{z_1z_2zy_n}$ are of one of the following three forms:

(i) $y_n^2y_jz_1$ (for some $j < n - 1$)
(ii) $y_n^2y_jy_k$ (for some $j, k < n - 1$)
(iii) $y_n^2z_1^2$.

In these three cases the $m = m' : M$ is

(i) $m = y_n^2y_jz_1 : z_1z_2zy_n = y_jy_n$,
(ii) $m = y_n^2y_jy_k : z_1z_2zy_n = y_jy_ky_n$, and
(iii) $m = y_n^2z_1^2 : z_1z_2zy_n = y_nz_1$ respectively.

However each of these are not minimal as $y_j, z_1 \in Q$ for $j < n - 1$.

So our ordering of our generators is a linear quotients ordering through the end of stage (2a).

4.1.3. *Stage (2b):* The second part of the second stage involves adding monomials $M = xy_iy_jz_2$ to our ideals I_M according to the lex order on (i, j).

$$\begin{aligned} Q &= I_{xy_iy_jz_2} : (xy_iy_jz_2) \\ &= (J^2 + (z_1z_2xy_j \mid 1 \le j \le n) + (xy_{i'}y_{j'}z_2 : (i', j') >_{\text{lex}} (i, j)) : (xy_iy_jz_2) \end{aligned}$$

Note the following inclusions hold, via the elements noted.

- $Q \supseteq (y_k \mid k < j)$ as $y_k = xy_iy_kz_2 : xy_iy_jz_2$
- $Q \supseteq (z_1)$ as $z_1 = xy_iz_1z_2 : xy_iy_jz_2$
- $Q \supseteq (z_2)$ when $j \ne n$ as $z_1 = y_iy_jz_2^2 : xy_iy_jz_2$
- $Q \supseteq (y_k \mid k > j + 1)$ as $y_k = y_iy_jy_kz_2 : xy_iy_jz_2$
- $Q \supseteq (y_{j+1})$ when $i \ne j$ as $y_{j+1} = y_iy_jy_{j+1}z_2 : xy_iy_jz_2$
- $Q \supseteq (y_j)$ when $i \le j - 2$ and $j \ne n$ as $y_j = y_iy_j^2z_2 : xy_iy_jz_2$

Taken together for $M = xy_iy_jz_2$ this gives

$$Q \supseteq \begin{cases} (y_1,\dots,y_n,z_1,z_2) & j \neq n, i < j-1 \\ (y_1,\dots,y_{j-1},y_{j+1},\dots,y_n,z_1,z_2) & j \neq n, i+1 = j \\ (y_1,\dots,y_{j-1},y_{j+2},\dots,y_n,z_1,z_2) & j \neq n, i = j \\ (y_1,\dots,y_{n-1},z_1) & j = n. \end{cases}$$

Assume $m \in Q$ is a minimal monomial generator that is not linear. That is $m = m' : xy_iy_jz_2$ for some m' before $xy_iy_jz_2$. As m is minimal, its support cannot contain any of the variables in Q. Also if x were to be in $\operatorname{supp}(m)$ then x^2 would divide m'. As no there is no such $m' \in I_M$ before $xy_iy_kz_2$, we have $x \not| m$. Thus the support of m satisfies

$$\operatorname{supp}(m) \subseteq \begin{cases} \emptyset & j \neq n, i < j-1 \\ \{y_j\} & j \neq n, i+1 = j \\ \{y_j, y_{j+1}\} & j \neq n, i = j \\ \{y_n, z_2\} & j = n \end{cases}$$

Case $(j \neq n, i < j-1)$**:** There is nothing to check as x does not divide m and all other variables are in Q.

Case $(j \neq n, i+1 = j)$**:** In this case m must be a power of y_j. As m is not linear, $y_j^2 | m$ and hence $y_j^3 | m'$. However none of the generators of $I(G)^2$ are divisible by y_j^3.

Case $(j \neq n, i = j)$**:** In this case $\operatorname{supp}(m) \subseteq \{y_j, y_{j+1}\}$. As m is not linear, we have one of the following must hold:

(i) $y_j^2 | m$
(ii) $y_jy_{j+1} | m$
(iii) $y_{j+1}^2 | m$.

In these three cases respectively we must then have

(i) $y_j^3 | m'$
(ii) $y_j^2y_{j+1} | m'$
(iii) $m' \in \{y_j^2y_{j+1}^2, y_jy_{j+1}^3, y_{j+1}^4, xy_jy_{j+1}^2, xy_{j+1}^3, z_2y_jy_{j+1}^2, z_2y_{j+1}^3, xz_2y_{j+1}^2\}$.

Case **(i)** cannot happen, as y_j^3 does not divide any generator of $I(G)^2$. Similarly, in case **(ii)**, $y_j^2y_{j+1} | m'$ which would require $y_jy_{j+1} \in I(G)$, which is not a generator of the edge ideal of the anticycle.

Finally, in case **(iii)** all degree 4 monomials divisible by y_{j+1}^2 have been enumerated as possible m'. None of these are generators of $I(G)^2$ except for $m' = xz_2y_{j+1}^2$. This however occurs later in our order.

Case $(j = n)$**:** In this case $\operatorname{supp}(m) \subseteq \{y_n, z_2\}$. As m is not linear, one of y_n^2, y_nz_2 and z_2^2 divide m. If y_n^2 or z_2^2 divide m then y_n^3 or z_2^3 divide m'. However no generator of $I(G)^2$ is divisible by a cube of a variable. If $y_nz_2 | m$ then $m' = y_n^2z_2^2$ which is not a generator of $I(G)^2$.

4.1.4. *Stage (2c):* Showing that this part of the ordering is a linear quotients ordering can be done using its symmetry with Stage (2b). We wish to show that all

Q such that

$$\begin{aligned} Q &= I_{xy_iy_jz_1} : (xy_iy_jz_1) \\ &= \Big(J^2 + \big(z_1z_2xy_j \mid 1 \le j \le n\big) + \big(xy_ky_lz_2 \mid 1 \le k \le l \le n, k < n\big) \\ &\qquad + \big(xy_ky_lz_1 \mid (k,l) <_{\text{lex}'} (i,j)\big)\Big) : (xy_iy_jz_1) \end{aligned}$$

are again generated by variables. We first show that Q' is generated by variables, for

$$Q' = \Big(J^2 + \big(z_1z_2xy_j \mid 1 \le j \le n\big) + \big(xy_ky_lz_1 \mid (k,l) <_{\text{lex}'} (i,j)\big)\Big) : (xy_iy_jz_1),$$

where the $<_{\text{lex}'}$ denotes the lex ordering on y_i with the variables in reverse order from the $<_{\text{lex}}$ used in Stage (2b).

Via symmetry with Stage (2b), this Q' must have linear quotients via an identical proof. From this, we see

$$Q' = \begin{cases} (y_1, \ldots, y_n, z_1, z_2) & j \ne n, j < i-1 \\ (y_1, \ldots, y_{j-1}, y_{j+1}, \ldots, y_n, z_1, z_2) & i \ne 1, j+1 = i \\ (y_1, \ldots, y_{j-1}, y_{j+2}, \ldots, y_n, z_1, z_2) & i \ne 1, i = j \\ (y_2, \ldots, y_n, z_2) & i = 1. \end{cases}$$

Clearly $Q' \subset Q$. We note that Q and Q' only differ by a colon ideal of the form

$$\big(xy_ky_lz_2 \mid 1 \le k \le l \le n, k < n\big) : (xy_iy_jz_1).$$

The generators of Q which are not in Q' are of the form $xy_ky_lz_2 : xy_iy_jz_1$ and hence all must divisible by z_2.

Since $z_2 \in Q'$ in all cases, we see that Q is generated by variables for all monomials M added in this stage.

4.1.5. *Stage (2d):* For the final case of Stage 2, we add all monomials in $J \cdot K$ of the form $m = xy_iy_jy_k$ ordered via lex with $y_1 > y_2 > \cdots y_n$. Our colon ideals then are of the form

$$\begin{aligned} Q &= I_{xy_iy_jy_k} : (xy_iy_jy_k) \\ &= \Big(J^2 + \big(z_1z_2xy_j \mid 1 \le j \le n\big) \\ &+ \big(xy_ky_lz_2 \mid 1 \le k \le l \le n, k < n\big) + \big(xy_ky_lz_1 \mid 1 \le k \le l \le n, 1 < l\big) \\ &+ \big(xy_{i'}y_{j'}y_{k'} \mid 1 \le i' \le j' \le k' \le n, i'+2 \le k', (i',j',k') >_{\text{lex}} (i,j,k)\big)\Big) : (xy_iy_jy_k). \end{aligned}$$

The last set of generators in $I_{xy_iy_jy_k}$ are given by

$$\big(xy_{i'}y_{j'}y_{k'} \mid 1 \le i' \le j' \le k' \le n, i'+2 \le k', (i',j',k') >_{\text{lex}} (i,j,k)\big)$$

as the variables can be arranged with indices i', j', k' in increasing order, but $i'+2 \le k'$ as at least one pair of $\{y_{i'}, y_{j'}, y_{k'}\}$ must be nonadjacent in the anticycle graph. This forces the given inequality.

Our colon ideals now satisfy the following inclusions, via the elements noted.

- $Q \supseteq (y_l \mid l < j)$ as $y_l = xy_iy_ly_k : xy_iy_jy_k$
- $Q \supseteq (z_2)$ as $z_2 = xy_iy_kz_2 : xy_iy_jy_k$

- $Q \supseteq (z_1)$ as $z_1 = xy_iy_kz_1 : xy_iy_jy_k$
- $Q \supseteq (y_l \mid l \geq j+2)$ as $y_l = y_iy_jy_ky_l : xy_iy_jy_k$
- $Q \supseteq (y_{j+1})$ when $i+1 \leq j$ and $j+2 \leq k$ as $y_{j+1} = y_iy_jy_{j+1}y_k : xy_iy_jy_k$
- $Q \supseteq (y_j)$ when $i+2 \leq j$ and $j+2 \leq k$ as $y_{j+1} = y_iy_j^2y_k : xy_iy_jy_k$.

Together this gives

$$Q \supseteq \begin{cases} (y_1, \ldots, y_{j-1}, y_{j+1}, \ldots, y_n, z_1, z_2) & i = j-1 \text{ and } j+2 \leq k \\ (y_1, \ldots, y_{j-1}, y_{j+2}, \ldots, y_n, z_1, z_2) & i = j \text{ or } j = k, k-1 \\ (y_1, \ldots, y_n, z_1, z_2) & \text{otherwise.} \end{cases}$$

Assume $m \in Q$ is a minimal monomial generator that is not linear. That is $m = m' : xy_iy_jy_k$ for some m' before $M = xy_iy_jy_k$. As m is minimal, its support cannot contain any of the variables in Q. Also if $x|m$ then $x^2|m'$. As this does not happen for any m' before $xy_iy_jy_k$, $x \notin \operatorname{supp}(m)$. Thus the support of m satisfies

$$\operatorname{supp}(m) \subseteq \begin{cases} \{y_j\} & i = j-1 \text{ and } j+2 \leq k \\ \{y_j, y_{j+1}\} & i = j \text{ or } j = k, k-1 \\ \emptyset & \text{otherwise.} \end{cases}$$

Case ($i = j-1$ and $j+2 \leq k$): In this case, m must be divisible only by y_j and cannot be linear. Thus $y_j^2|m$ and $y_j^3|m'$ which does not hold for any generator $m' \in I(G)^2$.

Case ($i = j$ or $j = k, k-1$): In this case, m has its support contained in $\{y_j, y_{j+1}\}$. As in the previous case, if the support of m contains $\{y_j\}$, we obtain a contradiction.

If the support of m contains $\{y_{j+1}\}$ and then m' must the product of y_{j+1}^2 and two of x, y_i, y_j, y_k. However, for this to be a generator of $I(G)^2$ the two chosen vertices must both be adjacent to y_j. If $i = j$, then $m'xy_{j+1}^2y_k$ is the only possibility, but this comes after $xy_j^2y_k$ in our ordering. If $j = k$ or $j = k-1$ then $m' = xy_iy_{j+1}^2$ is the only possibility. This again lies after $M = xy_iy_jy_k$ in the ordering.

Other Cases: In the other cases, the quotient contains all variables (except x, but there is no term divisible by x^2 which occurs prior to M in the ordering.) Hence, Q must be generated by linear terms.

4.1.6. *Stage (3a):* Now we move on to adding those terms in K^2, meaning monomials in $I(G)^2$ which came from pairs of edges xy_i and xy_j. Our colon ideals will be of the form:

$$\begin{aligned} Q &= I_{x^2y_iy_j} : (x^2y_iy_j) \\ &= \Big(J^2 + \big(z_1z_2xy_j \mid 1 \leq j \leq n\big) + \big(xy_ky_lz_2 \mid 1 \leq k \leq l \leq n, k < n\big) \\ &\quad + \big(xy_ky_lz_1 \mid 1 \leq k \leq l \leq n, 1 < l\big) + \big(xy_iy_jy_k \mid 1 \leq i \leq j \leq k \leq n, i+2 \leq k\big) \\ &\quad + \big(x^2y_ky_l \mid 1 \leq k \leq l \leq n, 1 < l, (k,l) >_{\text{lex}} (i,j)\big)\Big) : (x^2y_iy_j). \end{aligned}$$

These colon ideals satisfy the following inclusions via the elements noted.

- $Q \supseteq (y_1)$ when $j > 3$ as $y_1 = xy_1y_iy_j : x^2y_iy_j$
- $Q \supseteq (y_1)$ when $i > 1$ as $y_1 = x^2y_1y_i : x^2y_iy_j$

- $Q \supseteq (y_k \mid 1 < k < j)$ as $y_k = x^2y_iy_k : x^2y_iy_j$
- $Q \supseteq (y_k \mid i+2 \leq k \leq n)$ as $y_k = xy_iy_jy_k : x^2y_iy_j$
- $Q \supseteq (z_2)$ when $i \neq n$ as $z_2 = xy_iy_jz_2 : x^2y_iy_j$
- $Q \supseteq (z_1)$ when $j \neq 1$ as $z_1 = xy_iy_jz_1 : x^2y_iy_j$

Together this gives

$$Q \supseteq \begin{cases} (y_3, \ldots, y_n, z_1, z_2) & i = 1, j = 2 \\ (y_1, \ldots, y_n, z_1, z_2) & i+2 \leq j \\ (y_1, \ldots, y_{j-1}, y_{j+1}, \ldots, y_n, z_1, z_2) & 1 < i = j-1 \\ (y_1, \ldots, y_{j-1}, y_{j+2}, \ldots, y_n, z_1, z_2) & 1 < i = j < n \\ (y_1, \ldots, y_{n-1}, z_1) & i = j = n. \end{cases}$$

Assume $m \in Q$ is a minimal monomial generator that is not linear. That is $m = m' : x^2y_iy_j$ for some m' before $M = x^2y_iy_j$. Again, as m is minimal its support cannot contain any of the variables in Q. Also if $x|m$ then $x^3|m'$ which does not happen for any $m' \in I(G)^2$. Thus the support of m satisfies

$$\mathrm{supp}(m) \subseteq \begin{cases} \{y_1, y_2\} & i = 1, j = 2 \\ \emptyset & i+2 \leq j \\ \{y_j\} & i = j-1 \\ \{y_j, y_{j+1}\} & 1 < i = j < n \\ \{y_n, z_2\} & i = j = n. \end{cases}$$

We examine each of these cases individually.

Case $(i = 1, j = 2)$: In this case m is divisible by one of y_1^2, y_1y_2, y_2^2 and hence m' is divisible by $y_1^3, y_1^2y_2^2, y_2^3$. None of these can hold for m' a generator of $I(G)^2$.

Case $(i+2 \leq j)$: There is nothing to check as x does not divide m' and all other variables are in Q.

Case $(i = j-1)$: In this case m must be a power of y_j. As m is not linear, $y_j^2|m'$ and hence $y_j^3|m$. No generators of $I(G)^2$ are divisible by y_j^3 (or any third power of a variable.)

Case $(1 < i = j < n)$: In this case m is divisible by one of y_j^2, y_jy_{j+1} or y_{j+1}^2. If m' is to appear before $x^2y_iy_j$ in our list, it cannot be $x^2y_j^2, x^2y_jy_{j+1}$, nor $x^2y_{j+1}^2$. As $i = j$, the remaining possibilities for m are $xy_j^3, xy_j^2y_{j+1}, xy_jy_{j+1}^2$ or a monomial of degree four in y_j and y_{j+1}. However, none of these are generators of $I(G)^2$.

Case $(i = j = n)$: In this case m is divisible by one of y_n^2, y_nz_2, z_2^2. So m' is divisible by one of y_n^4, $y_n^3z_2$, z_2^2. There are no $m' \in I(G)^2$ such that the first two hold. For the last, if $z_2^2|m'$ and y_n does not divide m then m' must be one of $z_2^4, z_2^3x, z_2^3y_1, z_2^2x^2, z_2^2xy_n, z_2^2y_n^2$. None of these are in $I(G)^2$.

From this, we see that $I(G)^2$ has a linear quotients through Stage (3a).

4.1.7. *Stage (3b):* Finally, we add our generator $x^2y_1^2$ to our ideal $I_{x^2y_1^2}$. We only need to check that for this one remaining generator, the following colon ideal is generated by variables:

$$\begin{aligned} Q &= I_{x^2y_1^2} : (x^2y_1^2) \\ &= \Big(J^2 + \big(z_1z_2xy_j \mid 1 \leq j \leq n\big) + \big(xy_ky_lz_2 \mid 1 \leq k \leq l \leq n, k < n\big) \\ &+ \big(xy_ky_lz_1 \mid 1 \leq k \leq l \leq n, 1 < l\big) + \big(xy_iy_jy_k \mid 1 \leq i \leq j \leq k \leq n, i+2 \leq k\big) \\ &+ \big(x^2y_ky_l \mid 1 \leq k \leq l \leq n, 1 < l\big)\Big) : (x^2y_1^2). \end{aligned}$$

We have the following inclusions by the elements noted:

- $Q \supseteq (y_k \mid 1 < k \leq n)$ as $y_k = x^2y_1y_k : x^2y_1^2$
- $Q \supseteq (z_2)$ when $i \neq n$ as $z_2 = xy_1^2z_2 : x^2y_1^2$.

This gives us that our colon ideal satisfies $Q \supseteq (y_2, \dots, y_n, z_2)$.

So, if $m \in Q$ is a minimal non-linear monomial, then $\operatorname{supp}(m) \subseteq \{y_1, z_1\}$ and $m = m' : x^2y_1^2$ for some $m \in I(G)^2$ before $x^2y_1^2$. If $y_1|m$ then m' must be divisible by y_1^3. There is no such $m' \in I(G)^2$. Thus $\operatorname{supp}(m) = \{z_1\}$.

Since by assumption, m is not linear, $z_1^2|m$. Thus, $z_1^2|m'$ and the other variables dividing m' can only be z_1, x or y_1. There is no way to form a generator of $I(G)^2$ using only these variables as y_1 and x and z_1 are not adjacent to z_1^2. Hence, $Q = (y_2, \dots, y_n, z_2)$.

So this provides a linear quotients ordering on $I(G)^2$. □

5. Future Research

For higher powers of the edge ideal $I(A_n)^k$ of the anticycle, it is still unknown if all powers have a linear resolution, much less linear quotients. Construction of linear quotient orderings on $I(A_n)^k$ would accomplish this.

Question 5.1. Does $I(A_n)^k$ have linear quotients for $k \geq 3$?

We produced an ordering above on $I(A_n)^2$ by decomposing A_n into complementary subgraphs P_{n-1} and $A_n \setminus P_{n-1}$. While this order is nonunique, ordering the edges of $I(A_n)^2$ by decomposing the graph into the complementary subgraphs H and $G \setminus H$, then considering pairs of edges as appropriate, seems to produce linear quotients orderings with the clearest descriptions. Extending this order to $I(G)^k$ in a similar fashion has proven fairly difficult, even in the case of $I(G)^3$, but would be a natural next step after Theorem 4.2.

A problem of more general interest is to complete Theorem 1.3 of Herzog, Hibi and Zheng by answering the following question:

Question 5.2. Let G be the complement of a chordal graph. Does $I(G)^k$ have linear quotients for $k \geq 2$?

We might also ask for a description of all edge ideals whose powers eventually have linear resolutions.

Question 5.3. Can we exhibit classes of graphs G such that for all sufficiently large k,

(i) $I(G)^k$ has a linear resolution, or
(ii) $I(G)^k$ has linear quotients?

In [NP09], it was conjectured that graphs satistfying Question 5.3(i) are precisely those graphs G with a C_4-free complement. General conditions for the second class however remain open. It appears that anticycles A_n form such a class, but we wish to find more general conditions for the powers of an edge ideal of a graph to stabilize on linear quotients.

Acknowledgements. We would like to thank Irena Peeva and Eran Nevo for getting us interested in this topic and Adam Van Tuyl for many stimulating conversations. The first author would like to thank his advisor, Sara Faridi, for her direction and enouragement on this project. The second author would like to thank her advisor, Mike Stillman, for his detailed commentary on this paper, greatly improving the exposition.

References

[Con06] Aldo Conca, *Regularity jumps for powers of ideals*, Commutative algebra, Lect. Notes Pure Appl. Math., vol. 244, Chapman & Hall/CRC, Boca Raton, FL, 2006, pp. 21–32. MR 2184787 (2007c:13017)

[Frö90] Ralf Fröberg, *On Stanley-Reisner rings*, Topics in algebra, Part 2 (Warsaw, 1988), Banach Center Publ., vol. 26, PWN, Warsaw, 1990, pp. 57–70. MR 1171260 (93f:13009)

[HHZ04] Jürgen Herzog, Takayuki Hibi, and Xinxian Zheng, *Monomial ideals whose powers have a linear resolution*, Math. Scand. **95** (2004), no. 1, 23–32. MR 2091479 (2005f:13012)

[HVT07] Huy Tài Hà and Adam Van Tuyl, *Splittable ideals and the resolutions of monomial ideals*, J. Algebra **309** (2007), no. 1, 405–425. MR 2301246 (2008a:13016)

[HVT08] Huy Tài Hà and Adam Van Tuyl, *Monomial ideals, edge ideals of hypergraphs, and their graded betti numbers*, J. Algebraic Comb. **27** (2008), 215–245.

[MV10] Susan Morey and Rafael H. Villarreal, *Edge ideals: algebraic and combinatorial properties*, arXiv:1012.5329v3 [math.AC] (2010).

[Nev11] Eran Nevo, *Regularity of edge ideals of C_4-free graphs via the topology of the lcm-lattice*, J. Combin. Theory Ser. A **118** (2011), no. 2, 491–501. MR 2739498

[NP09] Eran Nevo and Irena Peeva, *Linear resolutions of powers of edge ideals*, preprint (2009).

[Vil90] Rafael H. Villarreal, *Cohen-Macaulay graphs*, Manuscripta Math. **66** (1990), no. 3, 277–293. MR 1031197 (91b:13031)

Andrew H. Hoefel, Department of Mathematics and Statistics, Dalhousie University, Halifax, Nova Scotia B3H 4R2
E-mail address: `andrew.hoefel@mathstat.dal.ca`

Gwyneth Whieldon, Department of Mathematics, Cornell University, Ithaca, NY 14850
E-mail address: `whieldon@math.cornell.edu`